

Exercises 4

1) Let $e \in \widehat{\text{End}}_{\text{nc}}(k)$ be an idempotent.

Show that if $P \triangleleft G$ is a finite p -subgroup then $\text{res}_P^G(e) = e|_P$ and that if P' is in the same component of $\Delta(G)/G$ as P (i.e. you can get from one to the other by a chain of inclusions and conjugations of non-trivial p -subgroups) then $\text{res}_{P'}^G(e) = \text{res}_{P'}^G(e)$.

Deduce that the idempotents corresponding to the components of $\Delta(G)/G$ are primitive.

2) Calculate $\widehat{\text{End}}_{C_p \times \mathbb{Z}}(k)$ and $\widehat{\text{Aut}}_{C_p \times \mathbb{Z}}(k)$. (Easy if you know the right spectral sequences, needs some work if not.)

3) Calculate $T(\text{SL}_2(\mathbb{Z}))$ at different primes ($\text{SL}_2(\mathbb{Z}) \cong C_6 *_{C_2} C_6$).

4) Calculate $T(C_p *_{C_p} C_{p^2})$.

5) There is an obvious surjection $C_4 *_{C_2} C_4 \rightarrow Q_8$. Calculate the inflation map $T(Q_8) \rightarrow T(C_4 *_{C_2} C_4)$.

6) Calculate $T(C_p \times \mathbb{Z})$ (it is an HNN extension).

What happened to the 1-dim representations of \mathbb{Z} ?

7) Calculate $T(C_p \times \mathbb{Z} \times \mathbb{Z})$. Is it finitely generated?

8) Calculate $T(\mathbb{Z}/p^\infty)$. (\mathbb{Z}/p^∞ means the p -torsion in \mathbb{Q}/\mathbb{Z} ; it acts on a graph with finite stabilisers).

9) For $G = A *_C B$, M a kA -module, N a kB -module such that $M|_C \cong N|_C$ stably, show that $M \cong M'$, $N \cong N'$ such that $M'|_C \cong N'|_C$, a genuine isomorphism.

Hint: Make M, N Gorenstein projective. Let F be a very big free module for G and set $M' = M \oplus F \perp_A$, $N' = N \oplus F \perp_B$. Use the Eilenberg trick.

10) Show that any stable automorphism $\phi: M \rightarrow M$ can be realised as a genuine automorphism $\phi': M' \rightarrow M'$.

Hint: take M Gorenstein projective. Find a projective

Ende contd

Moderately P big enough

(This is not strong enough for the construction $E(M; \mathcal{O})$ in general, but it suffices for Q6,7 above. You could try and formulate and prove the more general version).

11) Check that $\varphi \mapsto C(k, k; \varphi)$ is a group homomorphism (or use D).

Hint: use Exercises 3, Q6.

12) Show that if M is endotrivial then the natural map
 $M \otimes_k M^* \rightarrow \text{End}_k(M)$ is a stable isomorphism, hence $\text{End}_k(M) \simeq k$ stably.